

Holons on a meandering stripe

Oleg Tchernyshyov[†] and Leonid P. Pryadko

School of Natural Sciences, Institute for Advanced Study, Princeton, New Jersey 08540

(February 1, 2008)

To study a possible effect of transverse fluctuations of a stripe in a two-dimensional antiferromagnet on its charge dynamics, we identify elementary excitations of a weakly doped domain wall in the Hubbard model. Hartree-Fock numerics and analysis of fermion zero modes suggest that for $U \geq 3t$ charged excitations are mobile holons, $Q = 1$, $S = 0$. Each holon resides on a kink in the position of the domain wall. We construct a simple model in which transverse stripe dynamics is induced solely by motion of the holons. In the absence of spin excitations (spinons, $Q = 0$, $S = 1/2$), stripe fluctuations *do not* suppress a tendency to form a global charge-density order.

Physics of charge carriers confined to an antiphase domain wall in a two-dimensional antiferromagnet (AF) is one of the most important and least understood ingredients of the stripe^{1,2}-based approach³ to high- T_c superconductivity. Both the long-distance interactions^{3,4} and the short-distance physics^{5,6} seem to be important for understanding even the basic phenomenology of stripes. Given^{1,7-9} the existence of metallic stripes in an antiferromagnet, their physics should not be far different from that of a more conventional one-dimensional electron gas (1DEG). However, as suggested by model calculations,¹⁰ a metallic stripe with repulsive interactions placed in an AF environment may develop a spin gap. (In a conventional spin-rotation-invariant 1DEG a spin gap can only appear in the presence of attractive interactions.¹¹) With spin degrees of freedom gapped, coupling between adjacent stripes could lead either to a global superconducting order, or to an insulating charge-density wave (CDW) state.³ In a 1DEG with repulsive interactions, CDW correlations diverge more strongly,¹¹ and one would expect them to dominate superconductivity. A way around this difficulty was suggested¹² by Kivelson, Fradkin and Emery (KFE), who noticed that transverse fluctuations of conducting stripes would induce a phase mismatch between CDWs on neighboring stripes, effectively suppressing the interstripe coupling and the tendency to form a charge-ordered state. In contrast, such fluctuations *do not* suppress superconducting ordering, since the superconducting phase is not spatially modulated.

Transverse dynamics of insulating¹³ and metallic¹⁴ stripes has been previously studied by Zaanen *et al.* Of a particular interest to us is the example¹⁴ of a metallic stripe with a built-in coupling of charge motion and transverse fluctuations. The model describes an initially insulating stripe with a $4k_F$ CDW (1/2 electron per unit cell). Assuming that doped holes reside on rows adjacent to the site-centered stripe, Zaanen *et al.* found charge-1/2 solitons, which shift the domain wall one lattice constant sideways (left/right), thus acquiring a transverse flavor.

In this paper, we construct a similar model to study the effect of transverse fluctuations of a stripe on its charge dynamics. First, we employ Hartree-Fock (HF) numerics and an analysis of fermion zero modes to iden-

tify elementary excitations of a lightly doped domain wall in the Hubbard model. Of course, such a texture is an excited, only locally stable configuration within the HF approximation. Nevertheless, it could be a good starting point for understanding the nature of globally stable metallic stripes. We find that a weakly-doped domain wall is *bond-centered*. At moderate and strong coupling, $U \geq 3t$, holes doped into such a wall create kinks in the transverse position of the stripe (wiggles) and form mobile solitons with spin $S = 0$ and charge $Q = 1$ (holons, Fig. 1). Unlike their 1D counterparts, holons on a domain wall have a transverse degree of freedom, an isospin: a right (left) wiggle is ascribed isospin $\rho = +1/2$ ($-1/2$). In this respect, they resemble the solitons of Zaanen *et al.*¹⁴ Second, we construct an effective 1D model of a stripe whose transverse fluctuations are associated solely with the motion of the holons, while spinons (kinks of a similar nature but with $Q = 0$, $S = 1/2$) are frozen out. In contrast with the KFE arguments,¹² transverse fluctuations of a stripe in our model *do not* break phase coherence of CDWs on neighboring stripes. Therefore, they *do not* suppress the tendency to global CDW ordering, although the effect may be restored with the inclusion of spinons.




FIG. 1. Spin-charge separation on a bond-centered domain wall (dashed line) in an easy-axis AF ($J_z \gg J_x, t$). A hole decays into a spinon ($Q = 0$, $S = 1/2$) and a holon ($Q = 1$, $S = 0$) with isospins as shown (see text). Open symbols denote y -integrated spin and charge.

Numerical mean-field solutions. We have solved self-consistently Hartree-Fock equations for a linearly polarized AF, $\langle s_x \rangle = \langle s_y \rangle = 0$, $\langle s_z \rangle \neq 0$, for small and intermediate interaction strengths $U = 2 \dots 4t$ of the Hubbard model. At half-filling, we have found locally stable self-consistent solutions with a bond-centered domain wall. A site-centered wall has a slightly higher energy. A reduction of staggered magnetization on the wall induces

a 1D band in the middle of the Hubbard gap Δ (Fig. 2). Fermions confined to a site-centered wall have a Dirac spectrum $E = \pm v|k_x - \pi/2|$ and acquire a small mass (gap) $\tilde{\Delta} < \Delta$ in the case of a bond-centered wall. This gap arises because an electron moving along the wall feels a nonzero x -staggered magnetization (vanishing by symmetry for a site-centered wall). At half-filling, the Fermi energy lies between the two midgap bands ($E_F = U/2$).




FIG. 2. Left: midgap one-particle spectrum $E(k_x)$ of an AF with a straight domain wall. Self-consistent solution of Hartree-Fock equations. $U = 2.5t$, 48×27 sites. Dashed line: low-energy approximation, Eq. (5). States outside the gap are not shown for the site-centered stripe. Right: 4 Fermi patches in the Brillouin zone, Eq. (2), shown with eigenvalues (τ_3^x, τ_3^y) . Dashed line: Fermi surface of the noninteracting system.

A bond-centered domain wall *with a wiggle* is similar to an AF chain with a kink in x -staggered magnetization (Fig. 3). As in the model of polyacetylene,¹⁵ a pair of degenerate localized states (zero modes) appears in the middle of the smaller gap. A wall with a kink doped with a single hole has a $S = 0$, $Q = 1$ soliton at the wiggle (Fig. 4). In view of translational invariance, the charged defect can move along the wall. We thus identify it with a holon of the large- J_z limit (Fig. 1). At $U = 2 \dots 3t$ (rather weak coupling), holons are cigar-shaped like spin bags of Schrieffer *et al.*,¹⁶ a consequence of the Fermi surface nesting. Orientation of a holon along one or the other lattice diagonal is correlated with the direction of the wiggle (the transverse flavor). We have verified that holons are the lowest-energy states of doped charges on an empty wall for $U \geq 3t$ (at the HF level). At a weaker coupling, they bind into bipolarons with $Q = 2$, $S = 0$, and zero isospin.



FIG. 3. Left: Bond-centered stripe with a wiggle as a superposition of a site-centered domain wall (black arrows) and a 1D AF chain with a kink (open arrows). Right: Holons with alternating isospins form a site-centered stripe.




FIG. 4. Staggered magnetization $m(\mathbf{r}) = (-1)^{x+y} \langle s_z(\mathbf{r}) \rangle$ and hole density $1 - n(\mathbf{r})$ in a Hartree-Fock calculation at $U = 3t$, 24×24 sites. A bond-centered wall with 2 wiggles and 2 doped holes.

Continuum Hartree-Fock approximation. HF equations for a linearly polarized Hubbard AF read

$$-t \sum_{\langle \mathbf{r} \mathbf{r}' \rangle} \psi_{-s}(\mathbf{r}') + U[\delta n(\mathbf{r}) - s m(\mathbf{r})] \psi_s(\mathbf{r}) = E \psi_s(\mathbf{r}), \quad (1)$$

where $m(\mathbf{r}) = (-1)^{x+y} \langle s_z(\mathbf{r}) \rangle$ is the staggered magnetization and s is the *staggered* spin index. In what follows, density fluctuations $\delta n(\mathbf{r})$ will be neglected¹⁷ to restore charge-conjugation symmetry. Eq. (1) can be cast in a matrix form using Pauli matrices $\{\sigma_i\}$. Staggered spin s is an eigenvalue of σ_3 , the hopping term is proportional to σ_1 . Most importantly, zero modes of Eq. (1) are invariant under charge conjugation $\psi(\mathbf{r}) \rightarrow \sigma_2 \psi^*(\mathbf{r})$. If such modes are present, the system contains solitons with $Q = 0$, $S = 1/2$ (spinons) or $Q = 1$, $S = 0$ (holons).¹⁸

We have determined (see, e.g., Fig. 2, left) that low-energy midgap states induced by an empty domain wall live at one of the four “Fermi patches” with lattice momenta $|k_x| \approx |k_y| \approx \pi/2$ (Fig. 2, right). Introducing smoothly varying amplitudes of a fermion wavefunction,

$$\psi_s(\mathbf{r}) \approx \sum_{\alpha=\pm 1} \sum_{\mu=\pm 1} \psi_{\alpha\mu s}(\mathbf{r}) e^{i\pi(\alpha x + \mu y)/2}, \quad (2)$$

adds two more indices, $\alpha = \text{sgn } k_x$ and $\mu = \text{sgn } k_y$. Accordingly, we preserve only those Fourier components of magnetization which connect the Fermi patches:

$$\langle s_z(\mathbf{r}) \rangle \approx \sum_{\alpha=0}^1 \sum_{\mu=0}^1 m_{\alpha\mu}(\mathbf{r}) e^{i\pi(\alpha x + \mu y)}.$$

With the new indices come two more mutually commuting sets of Pauli matrices, $\{\tau_i^x\}$ and $\{\tau_i^y\}$, $i = 1, 2, 3$. In terms of these, $\mathbf{k} \approx \frac{\pi}{2}(\tau_3^x, \tau_3^y)$, $(-1)^x = \tau_1^x$, $(-1)^y = \tau_1^y$. The HF Hamiltonian becomes

$$H_{\text{HF}} = -2ita \sigma_1 \tau_3^x \partial_x - 2ita \sigma_1 \tau_3^y \partial_y - U \sigma_3 m(\mathbf{r}), \quad (3)$$

$$m \equiv m_{11} + m_{01} \tau_1^x + m_{10} \tau_1^y + m_{00} \tau_1^x \tau_1^y.$$

Only $m_{11}(\mathbf{r})$, staggered magnetization proper, survives in the bulk inducing the Hubbard gap $\Delta = U|m_{11}(\infty)|$.

On a straight domain wall in the x direction, $m_{01} = m_{00} = 0$, while $m_{10} \neq 0$. The spectrum of midgap states depends on the symmetry of the wall. If the stripe is site-centered (bond-centered), wall fermions have a gapless (gapped) spectrum. The absence of a gap can be

ascertained by finding a zero mode at the momentum $p_x \equiv -i\partial_x = 0$:

$$\sigma_2 \frac{d\psi}{dy} = \frac{U}{2ta} [\tau_3^y m_{11}(y) + i\tau_2^y m_{10}(y)] \psi(y). \quad (4)$$

To zeroth order in m_{10} , solutions of Eq. (4) are eigenstates of σ_2 , τ_3^y and, e.g., τ_3^x (each zero mode comes from a single Fermi patch), giving a total of 8 linearly independent zero modes. As usual,¹⁹ only half of these solutions [those with $\sigma_2 \tau_3^y m_{11}(+\infty) < 0$] are localized on the wall, so that there are 4 zero modes. Remarkably, in addition to the usual twofold spin degeneracy, there is another spin-like degree of freedom, which will prove to be the transverse flavor. The origin of isospin (at weak coupling) is thus exposed: compared to a 1D chain, there are twice as many “Fermi points” on a straight domain wall in 2D — see Fig. 2, right.

The difference between one-particle spectra of site and bond-centered walls arises in the first order in m_{10} . Eq. (4) has four zero modes if m_{10} is an odd function of y , i.e., for a site-centered wall.²⁰ On a bond-centered wall, $m_{10}(y)$ is even and fermions have a gap (Fig. 2):

$$E(0) = \pm U \langle m_1 \rangle \equiv \pm U \int m_{10}(y) \psi^\dagger(y) \psi(y) dy.$$

The one-particle spectrum $E(p_x)$ on a straight site-centered stripe can be determined approximately by starting with $p_x = 0$ [Eq. (4)] and treating the first term in Eq. (3) perturbatively. In the limit $p_x \rightarrow 0$, states outside the main gap can be neglected, which reduces the Hilbert space to the four zero modes [Eq. (4)]. By using the degenerate perturbation theory, we find a Dirac spectrum (dashed lines in Fig. 2, left):

$$E(p_x) \sim \pm v p_x, \quad v = 2ta \langle \tau_1^y \rangle. \quad (5)$$

Domain-wall holons at $U \ll t$. As illustrated in Fig. 3, magnetization on a *bond-centered* wall with a wiggle can be obtained by superimposing $m(\mathbf{r})$ of a straight *site-centered* wall and that of a spin chain with a kink in x -staggered magnetization. Away from the wiggle, $m_{00}(\mathbf{r}) = m_{01}(\mathbf{r}) = 0$. To simplify the discussion, we will neglect these components altogether. Decompose $m(\mathbf{r})$ into an x -independent part and the rest:

$$m(\mathbf{r}) = m^{(0)}(y) + m^{(1)}(\mathbf{r}), \\ m^{(0)}(-y) = -m^{(0)}(y), \quad m^{(1)}(\pm\infty, -y) = m^{(1)}(\pm\infty, y).$$

The Hamiltonian (3) can now be split in two parts:

$$H_{\text{HF}}^{(0)} = -2ita \sigma_1 \tau_3^y \partial_y - U \sigma_3 [m_{11}^{(0)}(y) + m_{10}^{(0)}(y) \tau_1^y], \quad (6)$$

$$H_{\text{HF}}^{(1)} = -2ita \sigma_1 \tau_3^x \partial_x - U \sigma_3 [m_{11}^{(1)}(\mathbf{r}) + m_{10}^{(1)}(\mathbf{r}) \tau_1^y]. \quad (7)$$

As shown above, the “transverse part” (6) has 4 zero modes for each p_x . Within this Hilbert space, $H_{\text{HF}}^{(1)}$ describes right and left-moving fermions with spin, which see a staggered magnetization

$$\langle m_1(x) \rangle = \int dy u^\dagger(y) [m_{10}(\mathbf{r}) + m_{11}(\mathbf{r}) \tau_1^y] u(y),$$

where $u(y)$ is a zero mode (4) of Eq. (6). The midgap fermion band acquires a gap of its own, $\tilde{\Delta} = U |\langle m_1(\infty) \rangle| < \Delta$, with two zero modes (one for each spin) inside this smaller gap. “Longitudinal” wavefunctions of the two zero modes satisfy the equation

$$\sigma_2 \frac{d\psi(x)}{dx} = \frac{U}{2ta \langle \tau_1^y \rangle} \tau_3^x \langle m_1(x) \rangle \psi(x). \quad (8)$$

The existence of two holon flavors can now be deduced from Eqns. (4) and (8). The zero modes have a finite norm only if

$$\sigma_2 \tau_3^y m_{11}^{(0)}(+\infty) < 0, \quad \sigma_2 \tau_3^x \langle m_1(+\infty) \rangle / \langle \tau_1^y \rangle < 0.$$

It follows then that the product of eigenvalues

$$\tau_3^x \tau_3^y = \text{sgn}[m^{(0)}(y = +\infty) \langle m_1(x = +\infty) \rangle \langle \tau_1^y \rangle] \quad (9)$$

can be identified with the holon isospin 2ρ . This can be seen by extrapolating Eq. (9) to larger values of U , which reduces the size of holons. We have $\langle \tau_1^y \rangle = \langle (-1)^y \rangle = (-1)^{y_0}$, where y_0 is the row number of the chain in Fig. 3. According to Eq. (9), if $\tau_3^x \tau_3^y = +1$, spins on the chain and to the right (left) of the wiggle are an extension of the upper (lower) AF domain, as for the $\rho = +1/2$ wiggle in Fig. 3. Thus, $\rho = \tau_3^x \tau_3^y / 2$. This identification is consistent with numerical HF solutions (Fig. 4), where $\tau_3^x \tau_3^y = \text{sgn} k_x \text{sgn} k_y$ can be inferred from the orientation of a holon.

Effective holon Hamiltonian. Large- J_z cartoons (e.g., Figs. 1 and 3) suggest that any configuration of a domain wall is uniquely represented by an interface drawn through integer (if a hole is present) or half-integer (bond, no hole) lattice points. In the absence of overhangs, the imaginary-time interface dynamics due to spin exchanges and hole hops can be described by an SOS-type model. (Locality in time assumes that coupling to the bulk spin waves can be neglected.²¹) In the dilute limit, mobile single-particle excitations are spinons and holons, each equipped with an isospin 1/2. When spinons are frozen out,²¹ the only remaining excitations are holons, and the SOS model reduces to that of (iso)spin-1/2 fermions with the short-range Hamiltonian

$$H = \sum_i [(-\tilde{t} \psi_{i\rho}^\dagger \psi_{i+1\rho} + \tilde{J}_x s_i^+ s_{i+1}^- + \text{h.c.}) \\ + \tilde{U} n_{i\uparrow} n_{i\downarrow} + \tilde{V} n_i n_{i+1} + \tilde{J}_z s_i^z s_{i+1}^z]. \quad (10)$$

Here $n_i \equiv n_{i\uparrow} + n_{i\downarrow}$ is the holon-number operator, $n_{i\rho} \equiv \psi_{i\rho}^\dagger \psi_{i\rho}$, and isospin operators are $s_i^z \equiv (n_{i\uparrow} - n_{i\downarrow})/2$ and $s_i^\pm \equiv \psi_{i\uparrow}^\dagger \psi_{i\downarrow}$. Taking $\tilde{U} \rightarrow +\infty$ excludes double occupancies, while isospin exchange $\tilde{J}_x$ allows two holons on one side of the stripe to hop to the other side. By construction, we do not expect isospin-rotation invariance; in general, $\tilde{J}_z \neq \tilde{J}_x$.

Holons can be transported between different stripes without disturbing the AF order only in pairs (with opposite isospins). In contrast to the model of a quarter-filled chain¹⁴ with the same Hamiltonian (10), the field

$\psi^\dagger$ describes $Q = 1$ particles, and the appropriate pairing operator $\psi_{i\uparrow}^\dagger \psi_{i+1\downarrow}^\dagger$ has charge $Q = 2$. This operator is a combination of the (iso)singlet (SP) and an (iso)triplet (TP_0) pairing operators,

$$\mathcal{O}_{\text{SP}}, \mathcal{O}_{\text{TP}_0} = (\psi_{i\uparrow}^\dagger \psi_{i+1\downarrow}^\dagger \mp \psi_{i\downarrow}^\dagger \psi_{i+1\uparrow}^\dagger) / \sqrt{2},$$

and an instability in either of these channels could lead to global (Cooper-pair) superconductivity, even though isospin is local to a given stripe.

To analyze this scenario, we have performed a standard weak-coupling analysis^{11,22} at a generic density. Because a stripe horizontal on average is invariant under isospin reflection, isospin and charge degrees of freedom separate.²³ The scaling in each sector is determined by the usual constants K_s, K_c ; for repulsive interactions $K_s > K_c^{-1} > 1$. In the absence of the isospin gap, both SP and TP_0 correlation functions have the temperature exponent²² $\mu_{\text{SP}} = \mu_{\text{TP}_0} = K_c^{-1} + K_s - 2 > 0$, i.e., there is no divergence in either channel. Unlike in a more conventional 1DEG,¹¹ where the spin-rotation invariance requires $J_z = J_x$ and another mechanism (e.g., the “spin-gap proximity effect”¹⁰) is needed to develop the spin gap, here an isospin gap arises naturally for $\tilde{J}_z > |\tilde{J}_x|$ (an easy-axis anisotropy). At low temperatures, such a system is in the Luther-Emery phase, the TP_0 component freezes out, while the isosinglet pairing exponent becomes $\mu'_{\text{SP}} = K_c^{-1} - 2$; it diverges for $K_c > 1/2$. Isospins are ordered at $T = 0$ and the stripe becomes, on average, site-centered (Fig. 3, right).

As usual, the more divergent ($\mu_{\text{CDW}} = K_c - 2 < \mu_{\text{SP}}$) CDW correlations compete with the superconducting order. In our holon-only model (10), transverse stripe fluctuations¹² *never* suppress this instability. CDWs on neighboring stripes remain phase-coherent not only in the isospin-ordered phase (where stripe fluctuations are suppressed by the isospin gap), but also in the presence of fully developed transverse fluctuations.

Large- J_z cartoons indicate that the length of a stripe seen by a hole increases in the presence of spinons (cf. Ref. 12). This is why we believe that in the full interface model (with both spinons and holons present, as in Fig. 1) the transverse fluctuations may reduce the coherence of CDWs on neighboring stripes. However, other properties of this model require a separate study.

Acknowledgments. The authors thank M. M. Fogler, S. A. Kivelson, F. Wilczek and J. Zaanen for illuminating discussions. This work has been supported by DOE Grant DE-FG02-90ER40542.

- ² Theoretical predictions of stripes in the cuprates date back to J. Zaanen and O. Gunnarsson, Phys. Rev. B **40**, 7391 (1989); D. Poilblanc and T. M. Rice, Phys. Rev. B **39**, 9749 (1989); H. J. Schulz, J. Phys. (Paris) **50**, 2833 (1989).
- ³ S. A. Kivelson and V. J. Emery, in *The Los Alamos Symposium-1993: Strongly correlated electronic materials*, edited by K. S. Bedell *et al.* (Addison-Wesley, Reading, 1994), p. 619; V. J. Emery, S. A. Kivelson, and J. M. Tranquada, cond-mat/9907228 (unpublished).
- ⁴ U. Low, V. J. Emery, K. Fabricius, and S. Kivelson, Phys. Rev. Lett. **72**, 1918 (1994).
- ⁵ J. Zaanen, J. Phys. Chem. Solids **59**, 1769 (1998).
- ⁶ L. P. Pryadko *et al.*, Phys. Rev. B (to be published); cond-mat/9905146.
- ⁷ C. Nayak and F. Wilczek, Phys. Rev. Lett. **78**, 2465 (1997).
- ⁸ S. R. White and D. J. Scalapino, Phys. Rev. Lett. **81**, 3227 (1998); cond-mat/9907375 (unpublished).
- ⁹ C. S. Hellberg and E. Manousakis, Phys. Rev. Lett. **83**, 132 (1999).
- ¹⁰ V. J. Emery, S. A. Kivelson, and O. Zachar, Phys. Rev. B **56**, 6120 (1997).
- ¹¹ V. J. Emery, in *Highly conducting one-dimensional solids*, edited by J. T. Devreese, R. P. Evrard, and V. E. van Doren (Plenum Press, New York, 1979), p. 247.
- ¹² S. A. Kivelson, E. Fradkin, and V. J. Emery, Nature **393**, 550 (1998).
- ¹³ H. Eskes *et al.*, Phys. Rev. B **58**, 6963 (1998).
- ¹⁴ J. Zaanen, O. Y. Osman, and W. van Saarloos, cond-mat/9804300 (unpublished).
- ¹⁵ W. P. Su, J. R. Schrieffer, and A. J. Heeger, Phys. Rev. B **22**, 2099 (1980).
- ¹⁶ J. R. Schrieffer, X.-G. Wen, and S.-C. Zhang, Phys. Rev. Lett. **60**, 944 (1988).
- ¹⁷ As charge order is driven by spin order in this case, density fluctuations do not play a crucial role.
- ¹⁸ Our proof requires only invariance under “combined parity” $\psi(\mathbf{r}) \rightarrow \sigma_1 \psi(-\mathbf{r})$ and an energy gap separating zero modes from the rest of the spectrum.
- ¹⁹ See, e.g., R. Rajaraman, *Solitons and instantons: an introduction to solitons and instantons in quantum field theory* (North-Holland, New York, 1982).
- ²⁰ This actually holds to any order in m_{10} .
- ²¹ Magnons and spinons are suppressed in the presence of a spin gap (created, e.g., by an Ising anisotropy, $J_z > J_x$).
- ²² T. Giamarchi and H. J. Schulz, Phys. Rev. B **33**, 2066 (1986).
- ²³ A *tilted* domain wall is equivalent to a 1DEG in the presence of external polarizing magnetic field (Ref. 24), where spin and charge degree of freedom no longer decouple. Most importantly, backscattering averages out to zero and the isospin gap does not develop at a generic filling. Although even a small tilt modifies the scaling exponents μ_{SS} and μ_{CDW} , it cannot change their sign, and we conclude that, in the absence of spinons, a tilted stripe does not have a superconducting instability.
- ²⁴ T. Ogawa and H. Otani, J. Phys. Soc. Jap. **64**, 3664 (1995); T. Kimura, K. Kuroki, and H. Aoki, Phys. Rev. B **53**, 9572 (1998).

[†] Electronic address: oleg@sns.ias.edu.

¹ For a recent experimental review on stripe phases see J. M. Tranquada, J. Phys. Chem. Solids **59**, 2150 (1998).